

## Best practice in antimicrobial drug prescribing

**Clostridium difficile infection (CDI) is associated with antimicrobial use.  
Prescribing antimicrobials wisely can reduce the incidence.**

### Clostridium difficile infection (CDI)

- ❑ *C. difficile* is a bacterium present in the gut flora in some people.
- ❑ Antimicrobials disturb the balance of the gut flora, allowing *C. difficile* to multiply and cause infection.
- ❑ Symptoms of CDI can vary from mild diarrhoea to fatal bowel inflammation.
- ❑ *C. difficile* spores are shed in the faeces. The spores can survive for long periods in the environment. If ingested, they can transmit infection to others.

### Prudent antimicrobial prescribing

- ❑ Only prescribe antimicrobials when indicated by the clinical condition of the patient or the results of microbiological investigation.
- ❑ Do not prescribe antimicrobials for sore throat, coughs and colds in patients at low risk of complications.
- ❑ Consider delayed prescriptions in case symptoms worsen or become prolonged.
- ❑ If an antimicrobial is required, follow local guidelines.
- ❑ Choose a narrow-spectrum agent where possible and prescribe a short course.
- ❑ Generally, no more than 5-7 days' treatment is required.
- ❑ Three-day courses are appropriate in some cases.
- ❑ Broad-spectrum antimicrobials should be reserved for the treatment of serious infections when the pathogen is not known.

### CDI and primary care

- ❑ CDI has commonly been associated with hospital stay but it is being recognised that many cases originate in the community, due to indiscriminate use of antibiotics.
- ❑ Patients most at risk are the elderly, particularly if they have medical conditions and are in close contact with others, e.g. in a care home, residential treatment centre or hospital.

### Antimicrobials to avoid where possible

The antimicrobials most strongly associated with CDI are:

- ❑ Second and third generation cephalosporins: cefaclor, cefuroxime, cefixime and cefpodoxime are examples for oral use
- ❑ Clindamycin
- ❑ Quinolones (associated with the virulent 027 strain of *C. difficile*): ciprofloxacin, levofloxacin, moxifloxacin, ofloxacin, norfloxacin.
- ❑ Long courses of amoxicillin, ampicillin, co-amoxiclav or cephalixin.

### Antimicrobials to choose

- ❑ All antimicrobials are associated with CDI, but those with lower risk are trimethoprim, penicillin V, tetracyclines and aminoglycosides.
- ❑ If antimicrobials are required, prescribe a short course and follow the local formulary.
- ❑ Where therapy has failed or there are special circumstances, obtain advice from a local microbiologist.

**How we use antimicrobials affects the whole community.**

## Which patients are most at risk of CDI?

Patients are more at risk of CDI if they are:

- ❑ Elderly
- ❑ Suffering from severe underlying diseases
- ❑ Immunocompromised
- ❑ In an environment where they are in close contact with one another (e.g. in a care home), particularly if hygiene is lacking.

Other factors that increase the risk of CDI are:

- ❑ Use of antimicrobials
- ❑ Recent gastrointestinal procedures
- ❑ Presence of a nasogastric tube

The use of proton pump inhibitors (PPIs) seems to increase the risk of CDI. Only prescribe PPIs when indicated.

## Reducing the risk of CDI

### Prudent antimicrobial prescribing

- ❑ Broad-spectrum antimicrobials are strongly associated with CDI.

### Isolating infected patients

- ❑ Isolating patients with CDI reduces the spread of infection in care homes and other places where people are in close contact with one another.

### Good hygiene

- ❑ Everyone should wash their hands with soap and water before and after each contact with a CDI-infected patient, including at home.
- ❑ Alcohol gel is effective against MRSA but not against *C. difficile* spores.
- ❑ The National Patient Safety Agency's Clean Your Hands campaign has been rolled out to primary care - see [www.npsa.nhs.uk/cleanyourhands](http://www.npsa.nhs.uk/cleanyourhands).
- ❑ Carers of CDI-infected patients should wear gloves and aprons.

Refer to the PCT's *Clostridium difficile* guidelines.

## When can broad-spectrum antibiotics be recommended?

There are few indications for broad-spectrum cephalosporins or quinolones in primary care. The following situations are the only indications for their first-line use. Refer to the PCT antimicrobial guideline.

Acute pyelonephritis	Ciprofloxacin, trimethoprim or co-amoxiclav
Acute prostatitis	Ciprofloxacin or trimethoprim
Simple gonorrhoea	Cefixime
Animal bite or human bite	Co-amoxiclav
Suspected meningitis in penicillin-allergic patient	Cefotaxime

When using broad spectrum antimicrobials, counsel patients at risk to be alert for signs of CDI and to stop their antimicrobial and seek medical help if diarrhoea develops.

If prescribing antimicrobials to patients with a history of CDI, refer to the PCT's *Clostridium difficile* guidelines.

## Bottom line

**C**lindamycin and broad spectrum antimicrobials are associated with CDI.

**D**on't prescribe antimicrobials when they're not needed.

**I**f an antimicrobial is indicated, prescribe a short course of a narrow-spectrum agent at the appropriate dose, as outlined in the PCT antimicrobial formulary.

Supporting *Medicines Q&A* documents are available at [www.nelm.nhs.uk/en/NeLM-Area/Evidence/Medicines-Q--A/](http://www.nelm.nhs.uk/en/NeLM-Area/Evidence/Medicines-Q--A/)

Date of preparation: October 2008

[Which antimicrobials are implicated?](#)  
[Are acid-suppressant medicines a risk factor?](#)  
[Are probiotics useful?](#)  
[Detail aid \(including planning matrix\)](#)